

# **The Effect of Postcard Reminders on Vaccinations Among the Elderly: A Block-Randomized Experiment<sup>1</sup>**

Nuole Chen<sup>2</sup> Kris-Stella Trump<sup>3</sup> Stacy Hall<sup>4</sup> Quan Le<sup>5</sup>

*Accepted at Behavioural Public Policy*

*This version pre copy-edits.*

*Please use DOI when citing: <http://dx.doi.org/10.1017/bpp.2020.34>*

## **Abstract:**

Prior research in the behavioral sciences has demonstrated that reminders can be an effective tool for encouraging health-related behavior changes. This article extends that literature by reporting the outcome of a randomized control trial of mailed vaccination reminders. In addition to making a substantive contribution regarding the efficacy of mailed reminders, this article also makes a methodological contribution: it illustrates how researchers can study the causal impact of an intervention even when a pure parallel trial is not possible. In this study, the Louisiana Department of Health sent postcard reminders regarding four recommended vaccinations (influenza, tetanus, shingles, and pneumonia) to 208,867 senior residents of Louisiana. We used block randomization and a stepped wedge design to assess the efficacy of the intervention. Individuals were blocked by their prior vaccine record and randomized to receive the postcard in one of four consecutive months (October-January). The reminder postcard had an overall positive effect on vaccination rates. The statistically significant and substantively small increase in overall vaccination rates was driven by participants who received the postcard reminder early in the intervention period.

**Keywords:** Vaccination, Public health, Reminders, Health behavior

<sup>1</sup> The authors would like to thank Caglayan Baser, Ekrem Baser, Kelly Bidwell, Amira Boland, Jake Bowers, Pompa Debroy, Tony Fox, Christopher Grady, Michael Gusmano, Alice Iannantuoni, Jessica Leight, Elana Safran, conference discussants, anonymous reviewers, and team members at the Office of Evaluation Sciences in the General Services Administration, the Immunization Program Office in the Louisiana Office of Public Health, and the National Vaccine Program Office in the Department of Health and Human Services. The findings were previously presented at the annual conventions of the Midwest Political Science Association (2018), American Psychological Association (2018), the Association for Psychological Science (2019), and the American Society of Health Economists (2019).

<sup>2</sup> University of Illinois at Urbana-Champaign; Office of Evaluation Sciences, U.S. General Services Administration. Address correspondence to Nuole Chen at [nchen3@illinois.edu](mailto:nchen3@illinois.edu).

<sup>3</sup> University of Memphis.

<sup>4</sup> Louisiana Department of Health.

<sup>5</sup> Louisiana Department of Health.

## **Introduction**

In this article, we present the results of a randomized control trial, testing whether a postcard reminder increased senior citizens' uptake of recommended vaccines. Our intervention is oriented toward directly changing patient behavior (rather than attitudes). In other words, it falls in the category of behavior-oriented interventions which are noted as promising in systematic reviews such as Betsch et al (2015) and Brewer et al. (2017). Behavior-oriented interventions target one subset of the broader determinants of vaccine uptake; these broader determinants include vaccine confidence, constraints, and calculation (see Betsch et al 2018 for a review). Prior research on behavior-oriented interventions has indicated that reminders in particular can be an effective tool for changing health-related behaviors including vaccine uptake (Stone et al. 2002), though reminders are not uniformly successful (Bourgeois et al. 2008). Given hitherto promising but mixed results in this category of interventions, understanding the conditions under which behaviorally informed interventions such as mailed reminders are most effective is an ongoing area of research (Brewer et al. 2017).

This study makes three primary contributions. First, and most importantly, it contributes to the literature on vaccine uptake by analyzing the efficacy of a postcard reminder sent to senior citizens in southern United States during the 2017-18 influenza season. Second, we contribute to the behavioral insights literature by presenting the results of a study in which the intervention had several behaviorally informed features, such as simple messaging and a call to immediate action, and identifying contexts that may increase the probability of success for these types of interventions. Finally, we carry out our study

using methods that allow causal identification even in situations where a “pure” control group is not feasible. As such, our methods can speak to a broad audience of researchers operating in applied contexts, who for various reasons are unable to withhold treatment from any study participants but who still wish to rigorously estimate the causal effect of their intervention.

The study was carried out during the 2017-18 influenza season in Louisiana through a collaboration between the Office of Evaluation Sciences<sup>1</sup> at the US General Services Administration and the Louisiana Department of Health. We evaluated the effectiveness of a vaccination reminder postcard, sent to Louisiana residents ages 65-70. The postcard referenced four vaccines typically recommended for people in this age group: influenza, shingles, tetanus, and pneumonia. This postcard was sent during the 2017-18 fall-winter season to residents who were 1) in the target age bracket, and 2) listed as (over)due at least one of the four vaccines in the Louisiana Immunization Information System (IIS). Due to the preference to not disrupt an existing reminder program, we could not withhold the postcard from a control group of Louisianans. Additionally, logistical constraints meant that the postcards could not be sent out all at once, but rather needed to be staggered over a period of four months. We used the staggered nature of the mailings to randomly assign when eligible Louisianans received the postcard. The result was a stepped wedge research design, in which we could identify the causal effect of the postcard on the probability of receiving any of the four vaccinations during the study period.

We find that the postcard positively affected the average number of four recommended vaccines received by study participants. The treatment effect is substantively largest and statistically detectable among individuals who receive the postcard in October. In addition to these main findings, we explore the effect of the postcard reminder on each of the four types of vaccines and find that the postcard boosts uptake of the flu and shingles (but not pneumonia or tetanus) vaccines. We also explore whether sending the postcard reminder in October is effective because individuals who receive the postcard in October have the postcard for more days compared to the other months, or because the month of October is different. We find that the month of October may be particularly important for vaccination reminders. We discuss the implications of our findings for practitioners and use exploratory analyses to develop hypotheses for future research.

### **Encouraging Vaccination**

We study an intervention that was designed to increase compliance with the Centers for Disease Control and Prevention's (CDC) recommended schedule of vaccinations for 65-70-year-old United States residents. The CDC recommends that people in this age range should receive vaccinations against four illnesses: influenza, tetanus (Td/Tdap), pneumococcal pneumonia, and herpes zoster (shingles) (CDC 2018a). Actual vaccination rates for these four vaccines currently do not meet official immunization rate targets, as identified by the CDC as part of the Healthy People 2020 objectives. Specifically, Healthy People 2020 identified the following vaccine-specific goals: that 30 percent of adults over age 60 receive a zoster (shingles) vaccine, 90 percent of adults over age 65 receive a pneumococcal vaccine, and 90 percent of adults over age 65 receive a seasonal influenza vaccine (U.S. Department of Health and Human Services 2018). Rates of

compliance with these recommendations are generally under target, but vary significantly by geographic and demographic indicators. It is in this context of imperfect compliance with recommended vaccine schedules that we evaluate the effectiveness of a postcard reminder to receive one's recommended vaccinations. In the section below, we review the recommendations for these vaccinations in more detail, illustrating the public health relevance of this intervention.

The influenza (flu) vaccine reduces the risk of getting the seasonal flu; the effectiveness of the vaccine varies significantly from year to year, but one study estimated average effectiveness across years to be around 59 percent (measured as reduction in the risk of contracting the flu) (Osterholm 2012). The CDC's initial calculations of effectiveness for the 2017-2018 influenza vaccine estimate a 36 percent reduction in risk among the vaccinated compared to the unvaccinated, with a confidence interval of 27 percent to 44 percent (CDC 2018b). This estimated effectiveness is higher than was anticipated early in the 2017-18 flu season, which saw a more dangerous flu season than usual (McNeil 2018). Prior analyses have indicated that provision and uptake of the flu vaccine is typically cost-effective (Lee 2010).

Among adults in the United States, vaccination rates are particularly low for the influenza vaccine: an estimated 43 percent of adults and 59 percent of children received the vaccine during the 2016-2017 flu season (CDC 2017). The flu becomes more serious as people age; among the elderly, the flu is associated with increased risks of complications and mortality. Flu vaccine uptake among the elderly is higher than among adults ages 18-64:

an estimated 65 percent of adults over 65 years received the flu vaccine during the 2016-2017 flu season compared with an estimated 38 percent of adults aged 18-64 years (CDC 2017). However, flu vaccination rates continue to lag behind the recommended levels of 90 percent of adults over 65 and 80 percent of adults between 18-64 (U.S. Department of Health and Human Services 2018).

Elderly adults (over the age of 65) are, in addition to the annual flu vaccine, recommended to receive vaccines against shingles, pneumonia, and tetanus. The exact recommended sequence varies by the patient's individual vaccination history and risk factors (CDC 2018a), making it more difficult for patients to know when they are due a given vaccine. For example, the pneumococcal vaccine can protect against diseases including pneumonia, meningitis, and sepsis, and comes in two forms: 13-valent pneumococcal conjugate (PCV13) and 23-valent pneumococcal polysaccharide vaccine (PPSV23). The type and number of doses of vaccines recommended varies somewhat by age, previous vaccination history and underlying medical conditions (CDC 2015). While there was an increase in adults over 65 estimated to receive the pneumococcal vaccine from 60 percent in 2012 to 64 percent in 2015, vaccination rates still lie well below recommended rates of 90 percent (Williams 2017; U.S. Department of Health and Human Services 2018). Overall, given lower than recommended rates of uptake for these vaccines, there is an opportunity for behavioral interventions to help increase vaccination rates.

### *Reasons for Low Compliance*

The reasons for lower-than-recommended vaccination rates include practical barriers, such as cost and accessibility issues, as well as attitude-based reluctance and refusals which can arise due to perceived lack of efficacy or fear of side effects. These variables can jointly be termed vaccine hesitancy (MacDonald et al. 2015). Psychological models of vaccine hesitancy emphasize variables such as complacency, convenience, and calculation (MacDonald et al. 2015, Betsch et al. 2015). A recent contribution further suggests the addition of constraints and collective responsibility for a “5C” psychological model of vaccine acceptance (Betsch et al. 2018). These frameworks can be used to diagnose barriers to vaccine uptake, and to tailor policy solutions that address those barriers. For example, a recent review of flu vaccine uptake finds that hesitancy for this vaccine is affected by a broad range of factors, but especially confidence and complacency (Schmid et al. 2017).

The intervention studied in this paper was inspired by a decision-science approach; this type of intervention attempts to guide behavior “without trying to change beliefs and attitudes” (Chapman 2019). As such, it falls in the category of behavioral interventions, which in turn have been noted as particularly promising in two recent systematic reviews (Brewer et al. 2017 and Betsch et al. 2015). This type of intervention specifically aims to overcome factors such as convenience or complacency, while not directly addressing confidence. Even though this category of vaccine uptake interventions typically does not reduce practical barriers such as cost or distance, it can be effective at overcoming psychological barriers such as forgetfulness. Additionally, behavioral interventions tend

to be relatively low cost (as is the intervention studied here), which also speaks in their favor. We thus acknowledge the position of this specific intervention type as promising, yet limited in the type of barrier it addresses. We suggest an intervention like the one studied here may be especially interesting for practitioners as a potential complement to other interventions, as combined strategies to encourage vaccine uptake may be more effective than isolated approaches (Dini et al. 2018).

### *Identifying and addressing barriers to vaccine uptake in Louisiana*

The Louisiana Department of Health regularly runs initiatives to increase vaccine uptake in the state. In the 2016-17 flu season (the year immediately prior to the intervention evaluation presented here), an estimated 58 percent of Louisiana adults age 65 and over received the flu vaccine (the Louisiana State's Healthy People 2020 target is 70 percent and the national target is 90 percent). 73.1 percent of Louisiana adults age 65 and over had received at least one of two pneumococcal vaccines (the national Healthy People 2020 target is 90 percent).<sup>2</sup> As a result, the Louisiana Department of Health has ongoing work to increase vaccination rates in the state.

Barriers to vaccine uptake that the Louisiana Department of Health has identified as particularly relevant to their work include: lower than desired public awareness about the importance and safety of vaccinations, lower than desired rates of strong healthcare practitioner recommendations, and lower than desired access to vaccine providers (including issues with transportation to healthcare providers and community pharmacies). As discussed above, a postcard intervention is part of the behavioral intervention toolkit;



as such, it may be a good tool for increasing awareness (which is related to reducing complacency in the parlance of the 5C framework of vaccine hesitancy). While the postcard is not intended to reduce practical barriers such as cost of transportation, it includes a reference to medical providers' recommendation to get vaccinated (more detail provided below), and as such may address the lower than desired rates of strong healthcare provider recommendations. With these barriers in mind, the Louisiana Department of Health initially fielded a postcard intervention in the 2016-17 flu season, and repeated it (with the addition of a random assignment component as described in this paper) in the 2017-18 flu season.

The postcard was deployed during flu season in part because, with respect to the flu vaccine in particular, there is evidence that significantly more people intend to get a flu shot than actually do each year (Galarce et al. 2011). Areas such as this, where people hold intentions to act but there is a lack of follow-through, are a classic avenue in which timely reminders can be effective at changing behavior. As an additional practical consideration, more involved interventions can be significantly more resource-intensive: while it is possible to increase flu vaccine uptake rates through financial incentives and/or mandates (Pitts et al. 2014), this avenue may not always be practically feasible or ethical (Bronchetti et al. 2015). Similarly, general education campaigns can be expensive yet have mixed results, sometimes achieving substantial increases (Kimura et al. 2007), but sometimes not creating any change at all (Dey et al. 2001, Chamberlain et al. 2015). These considerations further supported the choice of a behavioral intervention strategy.

### *Reminders as Tools in Health Care*

As described above, the postcard addressed barriers to uptake that were identified locally as prominent concerns that a postcard may partially yet feasibly address. In this section, we further situate the chosen intervention in the literature on the efficacy of reminders as a tool in public health interventions.

Prior research gives us reasons to expect that reminders can increase compliance with recommended vaccine schedules. For example, many adults who get vaccinated are not highly informed about vaccines (Dube 2013); instead of actively demanding vaccines, they get vaccinated due to provider recommendations or due to social norms (perceptions that getting vaccinated is the common thing to do). For people in this category, reminders may be an effective way to convey recommendations, social norms, and increase the salience of the need to get vaccinated. In addition, Betsch et al. (2015) point out that since people with strong anti-vaccine beliefs are a relatively small minority, it may be most productive to focus scarce vaccine promotion resources on the larger group of complacent or forgetful individuals.

Empirical research has validated that reminder-based interventions can improve healthcare related compliance, including medication adherence and health promoting behaviors (Fenerty et al, 2012, Fry et al. 2009). Stone et al. (2002) show that reminder or recall initiatives are an effective way to increase compliance with vaccine schedules, probably because these initiatives successfully reduce rates of forgetfulness and complacency. In the specific context of flu vaccines, McCaul et al. (2002) find that

sending reminder letters to elderly individuals who do not get a flu shot in the previous year increase their uptake of the flu vaccine by 8.6 percentage points from a baseline of 20 percent, while Yokum et al. (2018) find that sending reminder letters to Medicare beneficiaries increase flu vaccine uptake by between 0.4 and 0.9 percentage points from a baseline of 25.9 percent. However, reminders are not always effective: Bourgeois (2008) finds that sending personalized reminders to employees with a baseline rate of flu vaccination at 19 percent results in a statistically insignificant increase in uptake (5 percentage points). The current study adds to this accumulating body of evidence.

### *Postcard Design*

While many variables can affect the effectiveness of reminders, one element that may be consequential is the content and design of the reminder. For example, prior research has shown that vaccine uptake is significantly affected by physician recommendations (Bratic et al. 2016), which was also a barrier identified by the Louisiana Department of Health as relevant in this situation. While the most effective recommendations are likely made by one's own doctor in person, it is possible that highlighting healthcare providers' recommendation more generally helps increase a reminder's effectiveness. The postcard circulated in this study includes a statement that the vaccine sequence is recommended by the CDC, as well as an exhortation to talk to one's healthcare provider about the sequence.

Another factor that may cause people to fall behind on their vaccination schedule could be failure to follow through on intentions. One way to increase follow-through is to ask

people to act immediately rather than putting off the intended action to an indeterminate future time (Milkman et al. 2011, Hagger & Luszczynska 2014). The postcard circulated in this study encourages the recipient to “call your healthcare provider today”; this may help increase follow-through on intentions to get vaccinated. By referencing the CDC and encouraging recipients to act immediately by calling their healthcare provider, the content and design of the card uses behavioral insights to increase the effectiveness of the postcard reminder.

### **The Setting**

During the 2017-18 flu season, for the second year in a row, the Louisiana Department of Health sent out a postcard reminder to Louisianans aged 65–70. An image of the postcard appears in Figure 1. The postcard mentions all four of the vaccines recommended by the CDC for 65–70-year-old people (influenza, pneumococcal, zoster, tetanus). The postcard was sent to individuals who showed as being (over)due at least one of these four vaccines in the Louisiana Immunization Information System (IIS, also referred to as the vaccine registry).<sup>3</sup> Because the sample of recipients was drawn from the IIS early in the flu season, the sample includes all 65–70-year-old Louisiana residents except those who had all of their vaccinations and had already received the 2017-18 season’s flu vaccine by 9 September 2017 (the date of finalizing the study sample).<sup>4</sup> The upper age limit was imposed due to the Louisiana Department of Health’s interest in minimizing the number of postcards accidentally sent to deceased residents (the IIS does not receive timely updates regarding deaths). The postcard was designed by Pfizer, who also funded

postcard distribution in both the 2016-17 flu season (no evaluation of the effort occurred that season) and in the 2017-18 flu season (under consideration here).<sup>5</sup>

Because the postcard initiative was ongoing and had in the past been sent to all Louisianans who met the inclusion criteria mentioned above, the Louisiana Department of Health preferred to not exclude any eligible residents from receiving the postcard. Additionally, logistical constraints required that the postcards be sent in batches rather than all at one time. The staggered nature of the intervention allowed us to randomly assign the timing of postcard receipt, thus overcoming the absence of a pure control group in this study.

[Figure 1 about here]

## **Experimental Design**

We employed a block-randomized stepped wedge design to evaluate the postcard reminder intervention. The stepped wedge design, also called a waitlist design, allows researchers to evaluate a program that requires all individuals to receive treatment (Gerber & Green 2012, Hussey & Hughes 2007, Brown & Lilford 2006). We used this design to identify the causal effect of the treatment in the context of our experiment, where every individual must receive the postcard reminder and join the treatment group by the end of the study period.<sup>6</sup> We additionally used block-randomization to increase the precision of our analysis. We blocked by individuals' vaccination histories, as individuals with different vaccination histories could react differently to the postcard.

### *Stepped Wedge Design*

In the stepped wedge design, individuals were randomly assigned a month in which they receive treatment; the months were October, November, and December of 2017, and January of 2018. We considered individuals randomized to the January 2018 month as the control group. In all analyses below, we excluded vaccinations that we received during and after January 2018 (at which point all participants had received the treatment). In the analyses, we compared the vaccination behavior of individuals assigned to receive the postcard in October, November, and December 2017 (respectively) to the vaccination behavior of individuals who were assigned to receive the postcard in January 2018. Table 1 visualizes the stepped wedge design.

[Table 1 about here]

### *Block-Randomization*

We blocked treatment assignment by vaccination history because the postcard reminder might have different effects depending on the individuals' prior vaccination behavior. The participants entered the study with different vaccination histories. Some participants might be up to date on all vaccinations except for the seasonal flu shot. Others might have not been up to date on their vaccinations for decades. Yet others might be up-to-date on some vaccinations and behind on others. Since Louisiana only sent postcard reminders to people who were (over)due at least one vaccination, individuals who were completely up to date on vaccinations were not part of the study pool.<sup>7</sup>

We expected that these different vaccination histories were informative about the individuals' propensity to take up vaccinations. For example, if individuals had not been up to date on vaccinations for decades, they might have made a conscious decision to not vaccinate, which would be unlikely to change as a consequence of our intervention. On the other hand, among those individuals who were up-to-date on some vaccinations but not others, we might find a higher proportion of individuals who were not opposed to vaccinations, but had simply forgotten they were due a vaccination or had not gotten around to acting on a general desire to get vaccinated. We might expect postcard reminders to work well for those individuals.

Because the recommendations for the pneumococcal vaccine depended on the individual's health history and because the vaccination itself was composed of several different shots, our data (which did not include individual health information beyond vaccination status) did not allow us to decide conclusively which individuals were overdue pneumococcal vaccines. Therefore, by necessity, we relied on prior vaccination records for shingles, tetanus, and influenza to block recipients based on vaccination history.<sup>8</sup>

Individuals in each block were randomly assigned, with equal probability, to receive the postcard in one of the four treatment months. The four blocks are defined as follows:

1. *Vaccinated*: individuals who are up-to-date on both tetanus and shingles vaccines and also received a flu shot in the flu season immediately prior to our experiment.

2. *Non-vaccinated*: individuals who are overdue on both tetanus and shingles vaccines and did not receive a flu shot in the year prior to our experiment.
3. *Under-vaccinated (all but flu)*: individuals who are up-to-date on tetanus and shingles vaccinations but did not receive a flu shot in the year prior to the experiment.
4. *Under-vaccinated (mixed)*: individuals who were overdue some combination of tetanus, shingles, and flu vaccines.

The block sizes were different, but within each block, individuals had an equal (0.25) probability to be assigned to each of the four months. Table 2 gives further detail on the blocks and numbers of individuals within each block. In total, approximately 50,000 individuals received the postcard reminder every month.

[Table 2 about here]

We hypothesize that the postcard reminder will increase vaccinations among individuals in each of the treatment groups (October, November, and December) compared to control individuals. This hypothesis was pre-registered with the U.S. General Services Administration's Office of Evaluation Sciences. In the results section below, we further investigate whether the postcard had a greater impact after participants had been exposed to it for a longer time period, and whether its impact varied by vaccine. These analyses were not pre-registered and should therefore be interpreted as exploratory and hypothesis-generating.



## **Data and Methods**

Our data were drawn from the Louisiana Immunization Information System from the Louisiana Department of Health. Our experimental pool included 208,867 elderly individuals between the ages of 65-70 in Louisiana who were (over)due at least one vaccination at the time of the beginning of the experiment, 9 September 2017.<sup>9</sup> Table 3 presents baseline descriptive statistics about vaccinations in the experimental pool. At the start of the experiment, more individuals had received a recent seasonal flu vaccine,<sup>10</sup> as compared to having received the non-seasonal tetanus and shingles vaccines.

### *Missing Data*

The Louisiana Immunization Information System (IIS) records the last vaccination date of each vaccine received by each resident. Vaccinations records are compiled and submitted to the IIS by doctors and pharmacists in Louisiana. However, reporting vaccination dates for adults is voluntary for care providers. This means that data in the IIS in September 2017 were incomplete; while a number of individuals might have received their vaccinations, their health care providers might not have reported the data into the IIS. For the purposes of our analysis, the absence of a vaccination record means one of two things: that the individual did not receive the vaccine, or that the individual received the vaccine but the health care provider did not report it.

The rates at which vaccination records were absent vary significantly by vaccine:

- Flu: 28 percent of individuals have no record of receiving a seasonal flu vaccine (in any year)
- Pneumonia: 74 percent of individuals have no record of receiving this vaccine

- Tetanus: 68 percent of individuals have no record of receiving this vaccine
- Shingles: 81 percent of individuals have no record of receiving this vaccine

In other words, missing data were present in both treatment and control conditions, but due to data limitations, we could not distinguish missing data from non-vaccination. We chose to err on the side of caution and treat all null vaccination records as if the individual did not receive the vaccination. This means that in the analyses below, we estimated the average treatment effect of the postcard on receiving a vaccine that was reported to the IIS. The null outcome included both individuals who did not receive the vaccine and individuals who received the vaccine the healthcare provider but did not report the vaccination to the IIS. This choice means that in the results reported below, we most likely underestimated the impact of postcards on vaccine uptake.

### *Measurement*

The outcome of interest is in the proportion of the four recommended vaccinations received by the end of the observation period. For example, if an individual received two out of four vaccines (e.g. flu and shingles) in the month before receiving the intervention, the individual's baseline outcome is 0.5. If the individual then received an overdue vaccine (e.g. tetanus) after receiving the postcard, the post-treatment outcome is 0.75. We describe effects in percentage point changes throughout the rest of the paper; the percentage point estimate refers to the change in the proportion of vaccines received. In addition to this joint analysis of all four vaccines, we also carried out post-hoc analyses of the results for each vaccine separately and for each treatment month separately.

## *Method*

We use a saturated regression (Lin 2013) to estimate the average treatment effect of the postcard reminder. The outcome is the proportion of vaccines out of four that an individual received between 9 September 2017 – 9 January 2018. The treatment indicator denotes whether an individual is in the October, November, or December treatment groups, or the January control group. We also include 1) the number of vaccines the individual received between 8 September 2016 – 8 September 2017 as a baseline indicator and 2) the block indicator for each individual. This regression adjusts for covariates by interacting mean-deviated covariates with the treatment indicators. Additionally, because the outcome includes multiple comparisons, significance levels per outcome are reported both with unadjusted p-values and after using the Holm-Bonferroni procedure to adjust p-values (Holm 1979).

## **Results**

The postcard reminder has a small but statistically significant effect on vaccination among recipients who received the postcard in October (see Table 4).<sup>11</sup> These results hold with and without adjusting p-values. The first set of results in Table 4 shows the average treatment effect for individuals who received the postcard in October. We estimate that by 9 January 2018, individuals in the October treatment group received 0.27 percentage points more vaccinations (563 additional vaccinations) when compared to individuals in the control group; a difference that is statistically significant. Individuals in the November group received 0.15 percentage points more vaccinations (313 vaccinations) than the control group, and individuals in the December group received 125 more

vaccinations than the control group; however, the vaccination rates in the November and December treatment groups are not statistically significantly different from the control group.

For the pre-registered hypothesis, we find that receiving the postcard in October caused higher vaccination rates. Receiving the postcard in November or December did not have a statistically significant impact on vaccine uptake, though the point estimates were positive. Going beyond the aggregate results for the pre-registered hypothesis, we can ask more fine-grained albeit post-hoc questions about the impact of the reminder postcard. For example, do we see these results because the postcard was particularly effective for some subset of the four recommended vaccinations? Also, do we see time-varying results because sending the postcard in the month of October is particularly effective, or because individuals in the October treatment group were exposed to treatment (i.e. had the card in their possession) for longer? In the next section, we explore questions about the type of vaccination and how treatment month vs. length of treatment exposure might affect vaccination rates. As mentioned above, we consider the studies in the next section exploratory; we did not pre-register these outcomes nor make multiple comparison adjustments to the p-values of the analyses.

### *Effects by Vaccine Type*

First, we explore the possibility that the postcard is particularly effective for some subset of the four recommended vaccines. To do this, we estimate the effect of the postcard on each recommended vaccine separately, and report the results in Table 5. We find that receiving the postcard in October significantly increases uptake of the flu and shingles

vaccines, with the flu vaccine seeing the largest increase. The null results for the November and December treatment groups persist for all four vaccines. This suggests that the observed overall increase in vaccination rates in the October treatment group is driven by increased flu vaccinations, and to a lesser extent by increased shingles vaccinations.

[Table 5 about here]

### *Over-Time Variation*

Both Table 4 and Table 5 show that individuals who receive the postcard reminder in October have the highest vaccination rates when compared to the control group. However, these analyses do not allow us to distinguish whether vaccination rates are higher in October because receiving a postcard reminder during the month of October is particularly effective, or because having the postcard reminder for 3 months instead of 2 months or 1 month is more effective.

While our study was not pre-registered to distinguish between the effect of receiving the postcard in October and the effect of having a postcard reminder for 3 months, we can nonetheless make use of the randomization process to look at this question through exploratory analyses. Figure 2 depicts the number of vaccinations received by study participants during the observation period. Each line represents either the control group or one of the monthly treatment groups. We see that there are overall seasonal trends in the rate of vaccination, in that individuals tend to vaccinate more in October than in November or December, regardless of their assigned treatment group. We also observe

weekly trends: individuals tend to vaccinate more frequently during weekdays instead of weekends. These trends cannot speak directly to the origins of the October-specific effect, but we can use further analysis to tease apart a cumulative treatment time effect from a seasonal or October-specific effect. In particular, if the number of months that an individual has a postcard is important, then we would expect individuals who had the postcard reminder for 21 days<sup>12</sup> to have similar treatment effects, regardless of the month in which they received the card. On the other hand, if the treatment effect in October is truly different than the treatment effects in November and December, then we should see a difference in the estimated treatment effect even when we restrict observation to 21 days after receiving the postcard.

[Figure 2 about here]

Figure 3 compares the proportion of individuals who received at least one of the four vaccinations within 21 days after the postcard reminder was mailed to their treatment group, against the behavior of the control group in the same time period. Note that each of these three comparisons is made against the January control group. This means that we use the January control group three times at three different time points in these comparisons.<sup>13</sup> If the postcard reminder is more effective due to the length of time that an individual is exposed to the postcard, then we would expect the difference between treatment and control groups in Figure 3 to be fairly similar in each of the three months. However, Figure 3 shows that the difference between treatment and control groups is largest in October, and then increasingly smaller in November and December. This

difference suggests that the postcard reminder was more effective in October, and that this was not due to an individual's exposure to the postcard for 3 months, but rather due to the card being particularly effective when sent during the month of October.

[Figure 3 about here]

Finally, we use regression to estimate the effect of the postcard 21 days after the postcard reminder was sent for the October, November, and December groups. Table 6 reports these exploratory analyses, which suggest that the effect of the postcard is larger and statistically significant in October, smaller but still statistically significant in November, and smallest in December. This result does not appear to be due to the length of time that the individual has had a postcard reminder in their possession. In the discussion section below, we return to this particular result and use it to suggest some hypotheses for future research.

[Table 6 about here]

## **Discussion**

Increasing vaccination rates among the elderly is an important goal for the Louisiana Department of Health and the CDC. This study shows that a simple and relatively low-cost<sup>14</sup> intervention - sending reminder postcards to elderly individuals who are (over)due at least one vaccination - can modestly increase vaccination rates.

We sent behaviorally informed postcard reminders to 208,867 residents of Louisiana aged 65-70, who were listed as (over)due at least one of four vaccinations in the Louisiana Immunization Information System (IIS). The postcard had a small but

detectable effect on rates of vaccination: at the end of the four-month trial period, the share of recommended vaccinations received was 0.27 percentage points higher in the treatment group compared to the control group; this translated to an additional 563 vaccinations. Further analysis of the data revealed that this treatment effect was primarily driven by additional flu and shingles vaccinations received by individuals who were sent the postcard in the month of October.

More broadly, our study serves as an example of applied research in settings where withholding a treatment from some study participants is not feasible. A stepped wedge design allowed us to rigorously evaluate the intervention while also allowing all participants to receive the treatment by the end of the study. Specifically, by staggering the postcard reminders such that they were sent out in different months, and by randomly assigning individuals to receive the card in different months, we were able to estimate the effect of sending out the postcard, and to compare the effect of the postcard across different months.

These results have practical applications, and also point toward new hypotheses. Practically, the study demonstrates that a reminder intervention sent by mail can be effective at changing vaccine uptake among an elderly population who are due one or more vaccinations. The success of the intervention suggests that at least some of the hypothesized barriers – awareness/complacency, and strong recommendations from healthcare providers – were slightly alleviated by the postcard. However, the modest effect size underscores the importance of considering cost-benefit calculations before



implementing a reminder program. In this study, we do not have the requisite data to carry out a cost-benefit analysis, but we would encourage such analyses to accompany reminder programs wherever feasible.

Additionally, the estimated effect in our study is lower than that reported in previous studies. Better understanding the conditions (demographics, season, delivery format, and frequency of the reminder) that may affect the efficacy of reminders is an important avenue of research, potentially especially suited for future meta-analyses of existing trials across multiple settings. The modest effect size also points to the potential relevance of a barrier that was identified as important in this population such as access to transportation, but not addressed by the intervention. It is plausible that the severity of this practical barrier reduced the efficacy of the behavioral intervention and increasing awareness may not matter if no transportation solutions are available. The modest success of the postcard thus simultaneously illustrates the promise and the limitations of behavioral interventions: they may be most useful in settings where practitioners have accurately identified behavioral barriers as a *primary* barrier. We encourage future researchers to carefully reflect on the context in which a behavioral intervention is considered; interaction effects between different barriers are likely to be important in determining the success of any given intervention.

This study also points to new hypotheses. We find that the postcard is effective for flu and shingles vaccinations (but not pneumonia or tetanus vaccinations) and only during the month of October. One reason for this finding could be a heightened awareness of

vaccines in October. October, of course, is a particularly active month for campaigns that encourage people, especially the elderly, to get the seasonal flu vaccine. The Louisiana Department of Health ran a roster of outreach programs, which included free flu vaccines as well as an infant immunization campaign, during this period. Our data support this interpretation up by showing that there are more vaccinations in October than in the other months.

This interpretation raises the possibility that reminders may be particularly effective if they are sent in the context of other related campaign activity, or during times of otherwise heightened public awareness of vaccination requirements. In other words, vaccination campaigns may have a “the sum is greater than the whole of its parts” quality, with different pieces of a campaign reinforcing each other to create and maintain the salience of vaccination among the public. The fact that the reminder is most effective for the flu vaccination in particular further reinforces the interpretation that external cues about the importance of flu vaccines may have interacted with our reminder.

While we are not aware of any other outreach campaigns for the shingles vaccine in Louisiana during this time period, we cannot rule out that the “spillover” effect that we observe for the shingles vaccine in October may be related to such campaigns, if they happened at the same time as our intervention. We encourage future researchers to consider, even if informally, the broader persuasion/outreach environment that their interventions take place in. To study how a postcard intervention could be affected by the persuasion environment, study designs with multiple sites that have different

persuasion environments but similar outreach programs may be particularly relevant, as this would allow the identification of interaction effects between outreach initiatives.

The heightened awareness of the flu after November 2017 could be another reason why October was the most effective month. The 2017-18 flu season was particularly deadly and contagious. The CDC reported that influenza-like illness activity began to increase in November 2017 (CDC 2019). Baton Rouge General Medical Center reported a 450 percent increase in positive flu tests compared to the same time in the previous flu season. Addressing this development, Louisiana Department of Health offered free rounds of flu vaccines due to the severity of the flu season (Lussier 2018). This postcard intervention may have had the biggest effect in October 2017 because the flu had not started to spread yet. Once the flu season was underway and more people learned about how deadly that year's strain was, it is possible people began to vaccinate regardless of receiving a postcard reminder. Therefore, the postcard could have had no effect after October because, by November, everyone in Louisiana was encouraged to vaccinate. While this expectation would be difficult to study in a one-off study, meta-analyses of year-on-year outreach effects, in conjunction with data on the severity of the flu seasons, may be able to speak to it. More modestly, information about the severity of the flu season could be added as a randomized component to interventions when appropriate, measuring whether such information “cancels out” the efficacy of other intervention language.

It is unclear why the spillover effect did not extend to pneumonia and tetanus vaccines, but one potential explanation is that the latter two vaccines have more complicated recommended uptake schedules, compared to the seasonal flu vaccine and the one-off shingles vaccine. Future studies could explore both the seasonal and potentially cumulative effects of reminders, and potential spillovers between different vaccinations. As we suggested for influenza outreach above, this type of research would likely require collaborations between researchers and institutions that run multiple interventions at once. Such collaborations would allow designs where several interventions are randomized at once, developing a systematic sense of the recipients' information environment. Alternatively, to study spillover effects between different vaccinations within one site, a study could randomly assign a reminder regarding multiple vaccines or a reminder regarding one vaccine. The between-conditions difference in vaccines received would measure spillover between different vaccinations. To study the cumulative effect of reminders, a study could compare first-time recipients of postcard reminders with returning recipients. Strategies for identifying first-time recipients could leverage age thresholds, changes in eligibility requirements, and moving from one region to another.

Finally, our study also has important limitations. The participants in this study were a particular subset of individuals: residents of Louisiana, ages 65-70, who may have been (over)due one of the four target vaccinations. Whether the results of this study are more broadly applicable - to other demographic groups, other forms of reminders, or other vaccinations - is a question for future research. We also recognize that some subset of our

participant pool likely received the postcard for the second year in a row; due to data limitations we are unable to explore whether this affected the efficacy of the intervention, but considerations of repeated exposure will be important in future research as feasible.

Another important limitation arises from the difficulties of using a state immunization information system as a source of data. The information system in Louisiana holds a comprehensive list of the target population, but because reporting on adult vaccinations is not mandatory, vaccine records are almost certainly an undercount of real vaccination rates. For our study, this means that our estimated effect sizes are likely biased downward, and that real effects may be larger. Future research with alternative data sources could address this shortcoming. Finally, the modest effect sizes in this study also point to the importance of considering behavioral interventions in their broader context: as discussed in the introduction, behavioral interventions are most effective for addressing psychological barriers, but are less obviously applicable to practical barriers such as cost and accessibility. We encourage future researchers to pay attention to the environments in which behavioral interventions may be more/less effective in causing behavior change.

In sum, in accordance with the predictions of behavioral science, we find that a postcard reminder can modestly boost vaccination uptake among recipients. By including a stepped wedge design and block randomization in our evaluation of the study, we were able to compare the effect of the intervention in different months and for specific vaccines. We found that this intervention appeared most effective early in the flu season

(in October), and that it boosted uptake of the flu and shingles (but not tetanus or pneumonia) vaccines.

<sup>1</sup> The Office of Evaluation Sciences is an interdisciplinary team in the U.S. General Services Administration that translates and tests evidence-based insights into concrete recommendations for how to improve government.

<sup>2</sup> 2016 immunization rates from the CDC's AdultVaxView data service at <https://www.cdc.gov/vaccines/imz-managers/coverage/adultvaxview/index.html>; accessed 18 February 2018. The CDC's data do not include state-specific estimates of compliance with the tetanus and zoster vaccines among Louisiana's elderly population. Note that compliance data in AdultVaxView are not derived from the Louisiana Information Immunization System (which we use for tracking vaccine uptake in this study); this explains the difference in recorded vaccine uptake in our data and in the CDC's data service.

<sup>3</sup> Individuals ages 65-70 were part of this study if they were missing any of the following: the most recent seasonal flu vaccine, the tetanus, shingles, or pneumonia (either PPSV or PCV 13) vaccines.

<sup>4</sup> The Louisiana Department of Health determined that the study at hand does not constitute human subjects research. No identifiable data was shared with any researchers outside the Louisiana Department of Health; the analysis was performed on completely anonymous data.

<sup>5</sup> Pfizer provides the funds for postcard distribution, but does not access the medical records of Louisianans in the Immunization Information System. Pfizer was aware of the evaluation effort, but was not substantively involved in the study design, data analysis, write-up, presentation, or dissemination of this article, and they had no pre-publication review rights.

<sup>6</sup> We note that because of the large sample in our study, we do not reuse the control group in this main analysis. Therefore, the experimental design is a stepped wedge design, but the analysis of the experiment is not that of a typical stepped wedge design.

<sup>7</sup> Because the study pool was defined based on a data draw executed on 9 September 2017, the only individuals who are excluded from the pool are those who had received vaccines against tetanus, zoster, pneumococcal pneumonia, and the 2017-18 seasonal flu shot by that date. Note that while pneumococcal vaccine was used as a basis for including participants in the study, it was not used as a basis for block-randomization; see main text for more detail.

<sup>8</sup> While all individuals are recommended to receive the pneumococcal vaccine after the age of 65, which of the two available vaccines they receive, and when, can vary based on medical history and prior vaccination records. Given the low uptake of this vaccine, we are interested in measuring the impact of the postcard on overall uptake of it. We trust we can measure this reasonably, because in expectation, most individuals who have not yet received this vaccine are in fact due to receive it. However, because of the complicated scheduling for this vaccine, we choose to not use it as a basis for block-randomization, given that we cannot establish with certainty whether any given individual is in fact overdue on it.

<sup>9</sup> No identifiable data were shared with the research team. Louisiana Department of Health handled all identifiable details; we worked with de-identified data that only included vaccination-relevant fields.

<sup>10</sup> Having received a recent flu vaccine was measured as having received at least one seasonal flu vaccine between 8 September 2016 and 8 September 2017. Due to data limitations, we were unable to further distinguish between people who received the 2016-17 flu vaccine, people who were early takers of the 2017-18 flu vaccine, and people who had received both.

<sup>11</sup> 356 individuals attrited from the study.

<sup>12</sup> We choose 21 days because that is the number of days between the date that the December group postcard reminder was sent out and the end of the study. By limiting this analysis to 21 days, we can consider outcomes from the October, November, and December groups.

<sup>13</sup> The October treatment group outcome is the proportion of individuals who had vaccinations between 17 October 2017 to 6 November 2017 out of all individuals in the October group; the October control group outcome is the proportion of individuals who had vaccinations between 17 October 2017 to 6 November

2017 out of all individuals in the control group. Similarly, the November treatment group outcome is the proportion of individuals who had vaccinations between 21 November 2017 to 11 December 2017 out of all individuals in the November group; the November control group outcome is the proportion of individuals who had vaccinations between 21 November 2017 to 11 December 2017 in the control group. Finally, the December treatment group outcome is the proportion of individuals who had vaccinations between 20 December 2017 and 9 January 2018 out of all individuals in the December group; the December control group outcome is the proportion of individuals who had vaccinations between 20 December 2017 and 9 January 2018 out of all individuals in the control group.

<sup>14</sup> Unfortunately, we are unable to run a cost-benefit analysis of this intervention, because the total costs of the intervention are unavailable to the research team.

## References:

Betsch, C., R. Bohm, and G. B. Chapman. 2015. "Using Behavioral Insights to Increase Vaccination Policy Effectiveness." *Policy Insights from the Behavioral and Brain Sciences* 2 (1), 61–73.

Betsch, Cornelia, Philipp Schmid, Dorothee Heinemeier, Lars Korn, Cindy Holtmann, and Robert Böhm. "Beyond Confidence: Development of a Measure Assessing the 5C Psychological Antecedents of Vaccination." *PLOS ONE* 13, no. 12 (December 7, 2018)

Bourgeois, Florence T., William W. Simons, Karen Olson, John S. Brownstein, and Kenneth D. Mandl. 2008. "Evaluation of Influenza Prevention in the Workplace Using a Personally Controlled Health Record: Randomized Controlled Trial." *Journal of Medical Internet Research* 10 (1), 1–12.

Bratic, Julia S, Elisabeth R Seyferth, and Joseph A Jr Bocchini. 2016. "Update on Barriers to Human Papillomavirus Vaccination and Effective Strategies to Promote Vaccine Acceptance." *Current Opinion in Pediatrics* 28 (3), 407–12.

Brewer, Noel T., Gretchen B. Chapman, Alexander J. Rothman, Julie Leask, and Allison Kempe. 2017. "Increasing Vaccination: Putting Psychological Science Into Action." *Psychological Science in the Public Interest* 18 (3), 149–207.

Bronchetti, Erin Todd, David B. Huffman, and Ellen Magenheimer. 2015. "Attention, Intentions, and Follow-through in Preventive Health Behavior: Field Experimental Evidence on Flu Vaccination." *Journal of Economic Behavior and Organization* 116, 270–91.

Brown, C. A., & Lilford, R. J. 2006. The stepped wedge trial design: a systematic review. *BMC medical research methodology*, 6 (1), 54.

Chamberlain, Allison T., Katherine Seib, Kevin A. Ault, Eli S. Rosenberg, Paula M. Frew, Marielysse Cortes, Ellen A S Whitney, Ruth L. Berkelman, Walter A. Orenstein, and Saad B. Omer. 2016. "Impact of a Multi-Component Antenatal Vaccine Promotion Package on Improving Knowledge, Attitudes and Beliefs about Influenza and Tdap Vaccination during Pregnancy." *Human Vaccines and Immunotherapeutics* 12 (8), 2017–24.

Centers for Disease Control (CDC) "Surveillance of Influenza Vaccination Coverage." October 15, 2013. Accessed at



[http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s\\_cid%3Dss6204a1\\_x](http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid%3Dss6204a1_x) on 03/24/2017.

Centers for Disease Control and Prevention. Pneumococcal Vaccine Timing for Adults. November 30, 2015. Accessed at <https://www.cdc.gov/vaccines/vpd/pneumo/downloads/pneumo-vaccine-timing.pdf> on 02/15/2018.

Centers for Disease Control and Prevention. September 28, 2017. Flu Vaccination Coverage, United States, 2016-17 Influenza Season. Accessed at <https://www.cdc.gov/flu/fluview/coverage-1617estimates.htm> on 02/15/2018.

Centers for Disease Control and Prevention. 2018a. Recommended Immunization Schedule for Adults Aged 19 Years or Older, United States. 2018. Accessed at <https://www.cdc.gov/vaccines/schedules/hcp/imz/adult.html> on 02/18/2018.

Centers for Disease Control and Prevention. 2018b. Interim Estimates of 2017-18 Seasonal Influenza Vaccine Effectiveness - United States, February 2018. February 15, 2018. *Weekly*: 67(6); 180-185. Accessed at <https://www.cdc.gov/mmwr/volumes/67/wr/mm6706a2.htm> on 02/16/2018.

Centers for Disease Control and Prevention. 2019. Summary of the 2017-2018 Influenza Season. Accessed at <https://www.cdc.gov/flu/about/season/flu-season-2017-2018.htm> on 02/24/2020.

Chapman, Gretchen B. “A Decision-Science Approach to Health-Behavior Change.” *Current Directions in Psychological Science* 28, no. 5 (October 1, 2019): 469–74.

Dey, P, S Halder, S Collins, L Benons, and C Woodman. 2001. “Promoting Uptake of Influenza Vaccination among Health Care Workers: A Randomized Controlled Trial.” *Journal of Public Health Medicine* 23 (4), 346–48.

Dini, Guglielmo, Alessandra Toletone, Laura Sticchi, Andrea Orsi, Nicola Luigi Bragazzi, and Paolo Durando. “Influenza Vaccination in Healthcare Workers: A Comprehensive Critical Appraisal of the Literature.” *Human Vaccines & Immunotherapeutics* 14, no. 3 (March 4, 2018): 772–89.

Dubé, Eve, Caroline Laberge, Maryse Guay, Paul Bramadat, Réal Roy, and Julie Bettinger. 2013. “Vaccine Hesitancy: An Overview.” *Human Vaccines and Immunotherapeutics* 9 (8), 1763–73.

Fenerty, Sarah D , et al. 2012. "The effect of reminder systems on patient's adherence to treatment." *Patient Preference and Adherence* 2012 (6), 127–135.

Fry, Jillian P, and Roni A Neff. 2009 "Periodic Prompts and Reminders in Health Promotion and Health Behavior Interventions: Systematic Review." *Journal of Medical Internet Research* 11.2: e16.

Galarce EM, Minsky S, Viswanath K. 2011. "Socioeconomic status, demographics, beliefs and A(H1N1) vaccine uptake in the United States." *Vaccine* 29 (32), 5284-9

Gerber, A. S., & Green, D. P. 2012. *Field experiments: Design, analysis, and interpretation*. WW Norton.

Hagger, M. S., & Luszczynska, A. 2014. Implementation intention and action planning Interventions in health contexts: State of the research and proposals for the way forward. *Applied Psychology: Health and Well-Being*. 6 (1), 1-47.

Holm, S., 1979. A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics*, 65-70.

Hussey, M. A., & Hughes, J. P. 2007. Design and analysis of stepped wedge cluster randomized trials. *Contemporary clinical trials*. 28 (2), 182-191.

Johnson, DR, Nichol, KL, Lipczynski, K. 2008. Barriers to adult immunization. *American Journal of Medicine*, 121 (7 SUPPL. 2)

Kimura, Akiko C., Christine N. Nguyen, Jeffrey I. Higa, Eric L. Hurwitz, and Duc J. Vugia. 2007. "The Effectiveness of Vaccine Day and Educational Interventions on Influenza Vaccine Coverage among Health Care Workers at Long-Term Care Facilities." *American Journal of Public Health* 97 (4), 684–90.

Lee, Bruce Y, Rachel R Bailey, Ann E Wiringa, Abena Afriyie, Angela R Wateska, Kenneth J Smith, and Richard K Zimmerman. 2010. "Economics of Employer-Sponsored Workplace Vaccination to Prevent Pandemic and Seasonal Influenza." *Vaccine* 28 (37), 5952–59.

Lin, Winston. 2013. "Agnostic notes on regression adjustments to experimental data: Reexamining Freedman's critique." *The Annals of Applied Statistics* 7 (1), 295-318.

Lussier, Charles. (2018, January 30). Flu season rages in Louisiana; officials estimate 1,000 deaths expected. *The Advocate*.  
[https://www.theadvocate.com/baton\\_rouge/news/article\\_ee28457e-061c-11e8-a58d-63d48fcd77d.html](https://www.theadvocate.com/baton_rouge/news/article_ee28457e-061c-11e8-a58d-63d48fcd77d.html)

MacDonald, N.E., 2015. Vaccine hesitancy: Definition, scope and determinants. *Vaccine*, 33(34), 4161-4164.

McCaul, Kevin D, Rebecca J Johnson, and Alexander J Rothman. 2002. "The Effects of Framing and Action Instructions on Whether Older Adults Obtain Flu Shots." *Health Psychology* 21 (6), 624–28.

McNeil, Donald G. "The Flu Vaccine Is Working Better Than Expected, CDC Finds" 18 February 2018. *The New York Times*. Accessed at  
<https://www.nytimes.com/2018/02/15/health/flu-vaccine-effectiveness.html> on 02/18/2018.

Milkman, Katherine L, John Beshears, James J Choi, David Laibson, and Brigitte C Madrian. 2011. "Using Implementation Intentions Prompts to Enhance Influenza Vaccination Rates." *Proceedings of the National Academy of Sciences of the United States of America* 108 (26), 10415–20.

Osterholm, Michael T., Nicholas S. Kelley, Alfred Sommer, and Edward A. Belongia. 2012. "Efficacy and Effectiveness of Influenza Vaccines: A Systematic Review and Meta-Analysis." *The Lancet Infectious Diseases* 12 (1): 36–44

Pitts, Samantha I., Nisa M. Maruthur, Kathryn R. Millar, Trish M. Perl, and Jodi Segal. 2014. "A Systematic Review of Mandatory Influenza Vaccination in Healthcare Personnel." *American Journal of Preventive Medicine* 47 (3), 330–40.

Schmid, P., Rauber, D., Betsch, C., Lidolt, G. and Denker, M.L., 2017. Barriers of influenza vaccination intention and behavior—a systematic review of influenza vaccine hesitancy, 2005–2016. *PloS one*, 12(1).

Stone, Erin G., Sally C. Morton, Marlies E. Hulscher, Margaret A. Maglione, Elizabeth A. Roth, Jeremy M. Grimshaw, Brian S. Mittman, Lisa V. Rubenstein, Laurence Z. Rubenstein, Paul G. Shekelle. 2002. "Interventions That Increase Use of Adult Immunization and Cancer Screening Services: A Meta-Analysis," *Annals of Internal Medicine* 136, 641-651.

U.S. Department of Health and Human Services (HHS), Office of Disease Prevention and Health Promotion. February 16, 2018. *Healthy People 2020*. Accessed at <https://www.healthypeople.gov/2020/topics-objectives/topic/immunization-and-infectious-diseases/objectives> on 02/16/2018.


Williams WW, Lu P, O'Halloran A, et al. 2017. "Surveillance of Vaccination Coverage among Adult Populations — United States, 2015." *MMWR Surveill Summ* 66 (SS-11), 1–28.

Yokum, David, Julie C. Lauffenburger, Roya Ghazinouri, and Niteesh K. Choudhry. 2018. "Letters Designed with Behavioural Science Increase Influenza Vaccination in Medicare Beneficiaries." *Nature Human Behaviour* 2 (10): 743-749.


## Figure 1: Postcard Reminder for Vaccinations

Showing inside of folded card on the top, and the outside of folded card on the bottom.


Recipient's address is printed in the blank space.



**Figure 2.** Daily vaccinations received by study participants during the course of the study, by treatment month.



**Figure 3.** Proportion of individuals who had vaccinations within 21 days of receiving the postcard reminder by treatment month.



**Table 1.** Stepped Wedge Design. “Control” indicates month in which a group is in the control condition. “Intervention” indicates month in which a group is in the treatment condition. n = number of individuals in each group, and as a result, the number of individuals who move from the control to treatment conditions each month. Groups are made up of each of the 4 blocks of vaccination “types.” Probability of treatment assignment is the same across blocks, though block sizes within treatment groups are different.

	September	October	November	December
<b>October Treatment Group</b> (n=52,217)	Control	Intervention		
<b>November Treatment Group</b> (n=52,216)	Control		Intervention	
<b>December Treatment Group</b> (n=52,217)	Control			Intervention
<b>January Control Group</b> (n=52,217)	Control			



**Table 2.** Blocking scheme by vaccination history. The blocks are different sizes (i.e. have different numbers of individuals). Treatment assignment is distributed evenly across each of the months within each block.

Block	Description of Block*
Vaccinator (n=7,657)	Received a recent flu vaccine (last 13 months); up to date on shingles, tetanus
Non-vaccinator (n=99,669)	Overdue on flu, shingles, tetanus
Partial vaccinator (all but flu) (n=5,457)	Up-to-date on shingles, tetanus, overdue on flu
Partial vaccinator (mix) (n=96,084)	Mix of up-to-date/overdue on flu, shingles, tetanus
*Pneumococcal vaccination histories were not part of the blocking scheme. Vaccinations for pneumococcal are included in the analysis and results.	

**Table 3.** Baseline Data - Vaccinations. N = 208,867. Dataset includes individuals without vaccination data. All percentages include individuals with no data.

<b>Vaccine</b>	<b># of Individuals</b>	<b>% of Individuals</b>
Received a recent flu vaccine (Last flu shot between 9/1/2016 - 9/30/2017)	68,259	33%
Up-to-date on TD/Tdap vaccine (Last TD/Tdap booster between 9/1/2007 - 9/30/2017)	44,413	21%
Up-to-date on Zoster vaccine (Received the shingles vaccine)	39,551	19%

**Table 4.** Effect of Postcard Reminder on Vaccination Rates among Elderly

	Vaccination Rates	P-values (unadjusted)	P-values (Holm adjustment)
Effect of Postcard Reminder in October Group	0.0027 (0.0008)	< 0.01	0.01
Effect of Postcard Reminder in November Group	0.0015 (0.0008)	0.06	0.76
Effect of Postcard Reminder in December Group	0.0005 (0.0008)	0.48	1.00
Constant	0.0859 (0.0001)	< 0.01	< 0.01
<i>Obs: 208,511</i> <i>Regression included baseline outcome and block indicator.</i>			

**Table 5.** Effect of Postcard Reminder on Vaccination Rates by Vaccine

	Flu Vaccine	Tetanus Vaccine	Pneumonia Vaccine	Shingles Vaccine
Effect (October)	0.0071*** (0.0026)	0.0010 (0.0007)	0.0007 (0.0010)	0.0016*** (0.0005)
Effect (November)	0.0039 (0.0026)	0.0005 (0.0007)	0.0007 (0.0010)	0.0004 (0.0005)
Effect (December)	0.0030 (0.0026)	0.0000 (0.0006)	-0.0008 (0.0010)	-0.0001 (0.0005)
Constant	0.2984*** (0.0018)	0.0111*** (0.0005)	0.0278*** (0.0007)	0.0064*** (0.0003)
<p><i>*p &lt; 0.1, **p &lt; 0.05, *** p &lt; 0.01</i>  <i>Obs: 208,511 per vaccine</i>  <i>Regressions included baseline outcome and block indicator.</i></p>				

**Table 6.** Effect of Postcard Reminder on Vaccination Rates within 21 Days after Release Date

	October	November	December
Effect of Postcard Reminder	0.0042*** (0.0017)	0.0024** (0.0010)	0.0005 (0.0006)
Constant	0.1845*** (00064)	0.0568*** (0.0038)	0.0181*** (0.0022)
<p>*<math>p &lt; 0.1</math>, **<math>p &lt; 0.05</math>, *** <math>p &lt; 0.01</math>  <i>October Obs: 104,260</i>  <i>November Obs: 104,236</i>  <i>December Obs: 104,267</i>  <i>Regressions included block indicators.</i></p>			